

Must Not Be Consumed 48H Prior to Myocardial Exam Ne Pas Prendre 48H Avant Épreuve Myocardique

Medications / Médicaments

- Fiorinal
- Gravergol I
- nstantine
- Lenoltec
- Megral
- Midol
- Nervine
- Norgesic
- Novo Gesic
- Novo Propoxyn
- Oradine
- Aggrenox
- Apo-Dipyridamole FC
- Asasantine
- Novo-Dipiradol
- Persantin
- Apo-Theo
- Novo-Theophyl
- Pulmophylline
- Quibron
- Slo-bid
- Theo-SR
- Théo-Bronc
- Théochron
- Theolair
- Théolair-SR
- Théophylline
- Uniphyl
- Choledyl
- Rouphylline
- Aminophylline
- Phyllocontin
- Pentoxifylline Trenta
- Triaminicin
- Trianal
- Triatec -8
- Tylenol No1
- Tylenol No2
- Tylenol No3 Migraine
- Viagra
- Levitra
- Cialis
- 222, 282, 292
- AC&C
- Anacin Extra
- Atasol-8,15,30
- Cafergot
- Darvon-N
- Dristan
- Ergodryl
- Excedrin Extra
- Exdol-8,15,30
- Tecnal c1/4,c12

Aliments / Foods

- Café / Coffee
- Thé / Tea
- Chocolat
- Cacao
- Bananes / Bananas
- Boissons gazeuses / Soft drinks
- Boissons énergisantes / Energy drinks
- Decaffeinated beverages are not allowed
- Boissons décaféinés ne sont pas autorisés